

NO EVIDENCE FOR A SYSTEMATIC FEII EMISSION LINE REDSHIFT IN TYPE 1 AGN

Jack W. Sulentic

Instituto de Astrofísica de Andalucía, CSIC, Spain

sulentic@iaa.es

Paola Marziani

INAF, Astronomical Observatory of Padova, Italy

paola.marziani@oapd.inaf.it

Sebastian Zamfir

University of Wisconsin, Stevens Point, USA

szamfir@uwsp.edu

and

Zachary A. Meadows

University of Wisconsin, Stevens Point, USA

Zachary.A.Meadows@uwsp.edu

ABSTRACT

We test the recent claim by Hu et al. (2008) that FeII emission in Type 1 AGN shows a systematic redshift relative to the local source rest frame and broad-line $H\beta$. We compile high s/n median composites using SDSS spectra from both the Hu et al. sample and our own sample of the 469 brightest DR5 spectra. Our composites are generated in bins of FWHM $H\beta$ and FeII strength as defined in our 4D Eigenvector 1 (4DE1) formalism. We find no evidence for a systematic FeII redshift and consistency with previous assumptions that FeII shift and width (FWHM) follow $H\beta$ shift and FWHM in virtually all sources. This result is consistent with the hypothesis that FeII emission (quasi-ubiquitous in type 1 sources) arises from a broad-line region with geometry and kinematics the same as that producing the Balmer lines.

1. Introduction

Spectra of Type 1 AGN show a diversity of broad and narrow emission lines that provide direct insights into the structure and kinematics of photoionized, and otherwise excited, gas in the vicinity of the putative central massive object. Broad emission lines, like much studied $H\beta$ (e.g., Zamfir et al. 2010, hereafter Z10), are thought to arise in or near an accretion disk acting as the fuel reservoir for the central supermassive black hole ($\log M \sim 7 - 9.5 M_\odot$). $H\beta$ shows a diversity of line widths as well as profile shifts and asymmetries (Sulentic et al. 2000a). Despite this diversity some systematics have emerged and are best highlighted via the concept of two Type 1 AGN populations (Sulentic et al. 2000b).

Population A show the smallest broad-line widths $\text{FWHM } H\beta = 1000\text{--}4000 \text{ km s}^{-1}$ and includes the Narrow line Seyfert 1 (NLSy1) sources ($\text{FWHM} < 2000 \text{ km s}^{-1}$). Pop. A $H\beta$ profiles are currently best fit by a single Lorentz function. Population B sources show $\text{FWHM } H\beta = 4000\text{--}12000 \text{ km s}^{-1}$ and require two Gaussians (one unshifted and one redshifted) for a reasonable profile description. “Broad-line” $H\beta$ profiles as narrow as $\text{FWHM} = 500 \text{ km s}^{-1}$ (Zhou et al. 2006) and as broad as $\text{FWHM} = 40000 \text{ km s}^{-1}$ (Wang et al. 2005) have been found. Pop. A is predominantly radio-quiet while pop. B involves a mix of radio-quiet and the majority of radio-loud quasars.

Broad- and narrow-line profile shifts are known and the phenomenology can be confusing. Narrow emission lines like $[\text{OIII}]\lambda 5007\text{\AA}$ are regarded as a reliable measure of the local quasar rest frame except in the case of “blue outliers”, usually found in sources with $\text{FWHM } H\beta = 1500\text{--}3500 \text{ km s}^{-1}$ and weak $[\text{OIII}]$ (Véron-Cetty et al. 2001; Zamanov et al. 2002; Marziani et al. 2003b; Boroson 2005). Blue outliers show $[\text{OIII}]$ blueshifts as large as $\sim 1000 \text{ km s}^{-1}$. No pop. B sources with blueshifted $[\text{OIII}]$ are known at low z (or luminosity). Careful use of $[\text{OIII}]$ and $H\beta$ narrow line as rest frame measures suggests that broad $H\beta$ in pop. A sources rarely shows a systematic red or blue shift above the FWHM profile level. A blueshifted component or asymmetry is observed in some extreme FeII strong pop. A sources (Marziani et al. 2010). Pop. B sources show more complex line shift properties. The $H\beta$ profile usually shows two components: 1) a “classical” broad component (BC; $\text{FWHM} = 4000 - 5000 \text{ km s}^{-1}$) with zero or small (red or blue) shift, and 2) a very broad (VBC; 10000 km s^{-1}) and redshifted ($\gtrsim 1000 \text{ km s}^{-1}$) component. Composites involving the 469 brightest SDSS-DR5 quasars¹ suggest that these two components represent the underlying stable structure of $H\beta$ in pop. B sources.

Broad FeII emission has been found in type 1 quasars since the era of photographic

¹We acknowledge the use of SDSS quasar spectra: <http://www.sdss.org/collaboration/credits.html>

spectroscopy in the '60s. FeII emission blends are almost ubiquitous in a sample of the brightest (usually highest s/n) SDSS quasars (Z10). Circumstantial evidence has accumulated supporting the assumption that FeII emission arises in or near the emitting clouds that produce other low ionization lines like $H\beta$ (see e.g., Boroson & Green 1992; Marziani et al. 2003a; Sulentic et al. 2006). FWHM FeII appears to correlate with FWHM $H\beta$ over the full range where FeII can be detected (FWHM=1000-12000 km s^{-1}). This can be clearly seen at low z by observing the shape (e.g., smoothness) of the FeII 4450-4700 Å blue blend (and the FeII multiplet 42 line at 5018 Å) near [OIII]λ5007Å. In pop. A sources the blend resolves into individual lines while it becomes much smoother in pop. B sources. Sources with the strongest FeII emission also show a weakening of $H\beta$ emission as expected if the latter is collisionally quenched in the same dense medium where strong FeII emission can be produced (Gaskell 1985).

2. A Systematic FeII Redshift?

Obviously systematic line shifts place important constraints on models for the geometry and kinematics of the broad line region. The most famous example involves a systematic blueshift of high ionization lines (e.g., CIV λ1549Å) relative to low ionization lines (e.g., Balmer) especially in pop. A sources (e.g., Gaskell 1982; Sulentic et al. 2007; Richards et al. 2011; Wang et al. 2011). Evidence was recently advanced (Hu et al. 2008, hereafter H08) for the existence of a *systematic* redshift of FeII relative to [OIII]λ5007 (and hence the Balmer lines) in a majority of Type 1 quasars. This result, along with a narrower estimated FeII line width, has been ascribed to FeII emission arising in a region with dynamics dominated by in-fall and located at larger radius than the region producing the bulk of $H\beta$. H08 argue that the amplitude of the shifts correlates inversely with source Eddington ratio ($L/L_{Edd} \equiv \eta$). Interpretations for such an FeII redshift have already appeared (Ferland et al. 2009; Shields et al. 2010; Boroson 2011) reflecting the potential importance of such a first-order kinematic signature. Having worked on line spectra and profile shifts for many years we were surprised by the H08 claims and decided to test the hypothesis of a systematic FeII redshift. Could we have missed it?

First let us consider what we know. Most pop. A quasars show relatively symmetric unshifted Lorentz-like $H\beta$ profiles with $\text{FWHM} < 4000 \text{ km s}^{-1}$. In our work using the brightest ($g < 17.5$ or $i < 17.5$; Zamfir et al. 2008) SDSS DR5 quasars we processed spectra for ~ 260 pop. A sources (from a sample of 469 quasars; Z10) and we found no evidence for a systematic shift of FeII lines relative to $H\beta$ or [OIII]λ5007. Such an FeII shift should be easiest to detect in the brightest pop. A SDSS spectra with narrowest broad-line profiles and strongest FeII

emission. It is immediately suspicious that more and larger FeII redshifts are claimed for pop. B sources. In only one pop. A source in our sample SDSS J0946+0139 do we find a large H β peak (90% intensity level) redshift of 1830 km s $^{-1}$. This source is similar to OQ208 (Marziani et al. 1993 and discussed in H08) which shows $\Delta v_r > 2000$ km s $^{-1}$. SDSS J0946 is the only pop. A source with a large FeII redshift in our Z10 sample (1/260). Z10 found 19 quasars with an H β peak (9/10 fractional intensity) blueshifted more than -320 km s $^{-1}$ and 4 sources with the peak redshift more than +320 km s $^{-1}$. The remaining 241 pop. A sources showed no significant H β peak shift (figure 8 of Z10). Best FeII template fits to these sources show no significant difference in centroid redshift between FeII and H β . There are two possible causes of small and spurious H β (or FeII) shifts: 1) host galaxy contamination and 2) blue outliers. Except in rare cases host galaxy contamination is unlikely to induce systematic redshifts with the amplitudes reported by H08.

Extreme blue outliers with [OIII] blueshifts in the range 400-1000 km s $^{-1}$ are rare and therefore cannot be the cause of the large and systematic shifts reported in H08. In fact H08 selection criteria rejected sources likely to be seriously affected by 1) or 2).

H08 chose 4000+ sources from SDSS DR5 with computed s/n ≥ 10 . Z10 also used DR5 where $\sim 94\%$ of sources show s/n ≥ 10 . Our sample was magnitude-limited with a slightly shallower redshift upper limit ($z=0.7$ instead of 0.8). Why do we reach different conclusions about FeII shifts? A big part of the answer could involve how s/n was computed. H08 compute s/n over the range 4430–5500Å. This procedure overestimates the quality of the data because it includes major emission lines in the computation. We compute s/n in the range 5600–5800Å, which is free of strong lines and represents as close as one can approach to an estimate of continuum s/n near H β . Using our range the H08 sample shows mean and median s/n values of 10.6 and 7.4, respectively; approx. 65% show s/n < 10 . We find that only 182 spectra of our bright sample are included in the H08's. The majority of the H08's sources are lower s/n than those in our sample.

3. SPECFIT Analysis

3.1. A2 and B1 Composite Spectra

One cannot estimate reliable FeII line shifts using individual SDSS spectra for sources fainter than about $g \sim 17-17.5$. In rough order of importance our studies indicate that the accuracy of FeII shift measures depends on: 1) FeII strength and FeII/H β profile widths, 2) spectral s/n and 3) if estimates depend heavily on fits to the 4430-4680 Å blend, strength of HeII 4686 emission. Typical individual spectra used by H08 *show* too low s/n to allow

convincing conclusions about FeII shift and width – typical parameter uncertainties for individual sources are much larger than the ones connected with our high s/n composites (for a typical A2 source with $s/n \approx 20$ uncertainties of shift estimates are larger than $\pm 1500 \text{ km s}^{-1}$). Individual source spectra with large quoted FeII redshift and s/n near the sample median were extracted from the H08 sample and SPECFIT modelled. Using an FeII template with fixed shifts ranging from zero up to the largest values quoted by H08, χ^2 cannot distinguish between zero and e.g., 1000 km s^{-1} redshift in the majority of the sources.

The best recourse is to generate high s/n composite spectra. H08 argue that one cannot confirm or refute the existence of a systematic FeII redshift using composite spectra because of the large dispersion of FWHM, shifts and flux values for both H β and FeII. This is likely true for composites generated from random subsamples of sources but not true if one generates composites over more limited ranges of parameter values. One can generate binned composites over limited ranges of FWHM H β and FeII strength using the 4DE1 formalism (Sulentic et al. 2002; Bachev et al. 2004; Sulentic et al. 2007; Z10). 4DE1 bins A2 (FWHM H β = 1000-4000 km s^{-1} , $0.5 \leq R_{\text{FeII}} \leq 1.0$) and B1 (FWHM H β = 4000 – 8000 km s^{-1} , $R_{\text{FeII}} \leq 0.5$) are of particular interest because they include the largest numbers of sources in random samples. SPECFIT analysis (Kriss 1994; details in Marziani et al. 2009) of an A2 median composite involving $n = 130$ Z10 sources ($s/n \approx 90$) gives a best-fit consistent with zero FeII redshift. The situation for the B1 composite ($n=131$ sources from Z10; $s/n \approx 110$) is less constraining because lines are broader and FeII weaker. Table 1 reports FeII template shifts and 2σ uncertainties for SPECFIT tests discussed in this paper. We also report peak shifts of H β BC extracted from the best SPECFIT solutions along with H β “core” shifts measured at the centroid of the line peak after H β NC subtraction. In no case do we find a significant shift between FeII and the rest frame or between FeII and H β . We also do not find any significant FeII shifts if we restrict to sources with L/L_{Edd} ratio $\eta \leq 0.1$ (H08 suggested the shifts might be largest for low L/L_{Edd} sources).

3.2. A2 Composites Involving Only Sources with Large FeII Shifts?

Since we find no evidence for systematic FeII redshifts in our Z10 bright quasar sample composites it is useful to generate FeII shift composites using the H08 sample. We generate them within the 4DE1 context thereby restricting the ranges of FWHM H β , FeII relative strength (and likely also FWHM FeII) for each composite. Since the distribution of FeII shifts shown in H08 is continuous we focus on the sources with largest quoted shift values. If these shifts are not confirmed then smaller shifts are even less likely to be real. We therefore focus on constructing median composites for all H08 sources falling in 4DE1 bins A2 and B1

with H08 FeII redshift estimates $\geq 1000 \text{ km s}^{-1}$ (figure 1). Two composites were constructed for each spectra bin: 1) one with no restriction on FeII width (H08 do not constrain FWHM FeII in their template fits so it is sometimes very different from FWHM $H\beta$) and 2) one with FeII width constrained to the FWHM range of $H\beta$ in a particular bin (i.e., $\leq 4000 \text{ km s}^{-1}$ for A2 and $4000 - 8000 \text{ km s}^{-1}$ for B1). Upper and lower panels of figure 1 show bins A2 (FWHM $H\beta < 4000 \text{ km s}^{-1}$) and B1 (FWHM $H\beta = 4000 - 8000 \text{ km s}^{-1}$), respectively ($n=156$ for bin A2 and $n=240$ for bin B1). The $s/n \sim 55-60$ for both composites. Spectra show best-fit SPECFIT models superimposed. The left and center panels involved FeII templates fixed to the best fit and 1500 km s^{-1} shifts, respectively. Our template prescription is described in Marziani et al. (2009)

Graphical results for the best-fits are shown in the right panels of figure 1. Fits were performed over the range $\approx 4470 - 5450 \text{ \AA}$, where FeII and continuum emission account for the total flux making it the safest region for normalized χ^2 computations. χ^2 values are shown for the range of adopted FeII shifts. In order to estimate confidence intervals we considered a set of fits with displacements $\Delta v_r = +500n$, for integer $n = 0..4$, along with the best fit and a few additional Δv_r cases in proximity to the minimum χ^2 . One can see a clear preference for zero or near-zero fits. The significance of χ^2 variations is described by F statistics appropriate for ratios of χ^2 values (Bevington 1969). Given the large number of degrees of freedoms in the sampling range ($4500 - 4630$, $5040 - 5090$, $5310 - 5360 \text{ \AA}$) any χ^2 differences between two fits become significant at a 95% confidence level if $\chi^2/\chi^2_{\min} \approx 1.24$. The χ^2 values indicate that zero shift and “best shift” values in table 1 are not significantly different. All fits involving shifts $\gtrsim 500 \text{ km s}^{-1}$ are statistically significant. The middle panel of figure 1 upper row demonstrates visually that the fit with $\Delta v_r = +1500 \text{ km s}^{-1}$ (and even more the fits with larger displacement) do not reproduce the observed FeII emission.

Both the residuals and χ^2 results rule out any systematic redshift for at least half of the H08 sample (pop. A). Note especially the fits to the two relatively isolated multiplet 42 FeII lines between $H\beta$ and [OIII] $\lambda 4959$ and on the red wing of [OIII] $\lambda 5007$. The redshifted fit fails to include the blue side of the $4450 - 4700$ blend and the red side is confused by the frequent presence of HeII $\lambda 4686$. The latter line is not mentioned in the H08 study leaving us to conclude that it was not included in their fits. It can certainly give the impression of a redshift of the FeII blue blend, which is the most useful FeII diagnostic in the optical spectra of low redshift quasars (the red FeII blend is frequently affected by coronal lines as well as MgII host galaxy absorption in lower redshift sources). Pioneering principal component analysis of the BQS survey (Boroson & Green 1992) found that HeII $\lambda 4686$ equivalent width anticorrelates with sources luminosity (it is Eigenvector 2). There is a tendency for the H08 sources with largest FeII redshifts to favor a smaller and lower ($\log L \sim 40.3 - 41.0$) range of source luminosity than those with near zero shifts ($\log L \sim 40.3 - 42.0$). Thus the effect

of HeII will tend to play a larger role in the sources where the largest FeII redshifts have been found. H08 show composite spectra for five bins of FeII redshift in their Figure 12. The three bins involving largest FeII redshift sources show a prominent HeII signature that, if not subtracted, will increase the apparent significance of any assumed FeII redshift. Only the bin involving sources with no FeII redshift (within the uncertainties) shows no evidence of HeII emission.

3.3. What about Population B Sources?

Figure 9a of H08 suggests that a larger fraction of quasars with $\text{FWHM H}\beta > 4000 \text{ km s}^{-1}$ (Population B) show large FeII redshifts. The lower panels of our Figure 1 show SPECFIT models superimposed on a B1 median composite constructed from all H08 sources with quoted FeII redshift greater than 1000 km s^{-1} . The situation is certainly more ambiguous than for the A2 composite. It is hard to identify individual FeII features. Lines are broad, FeII is weak and under these conditions there are serious limitations on the reliability of FWHM and shift estimates for FeII (cf. Fig. 3 of Marziani et al. 2003a). The same analysis as done for A2 composite shows much poorer constraints on the FeII shift. The best fit yields $\Delta v_r \approx 760 \text{ km s}^{-1}$ but is not distinguishable from a zero shift solution. If one actually computes χ^2 values over the ranges $4474 - 4640 \text{ \AA}$, $5040 - 5105 \text{ \AA}$, $5320 - 5400 \text{ \AA}$, the χ^2 monotonically increases from 0 shift (figure 1, lower rightmost panel), although the increase remains insignificant until $\approx 1100 \text{ km s}^{-1}$, where $\chi^2/\chi_{\min}^2 \approx 1.21$. B1 FeII is too faint and the lines are too broad to make inferences about line shifts and widths. The claim of large FeII shifts are not, and cannot be, confirmed.

Kovačević et al. (2010) recently report an FeII study of SDSS quasars and any FeII redshifts they measure (their figure 16) are much smaller than those reported by H08 (the average FeII shift relative to the narrow lines is $100 \pm 240 \text{ km s}^{-1}$).

Returning to our previous list of major sources of uncertainty for FeII shift and FWHM estimates leads us to suggest that low spectral s/n and above average HeII strength are the culprits. The fit to the $4430 - 4680$ blue blend drives the best fit χ^2 results. The exclusion of HeII4686 from the H08 fits likely results in a tendency for HeII to “redshift” the blue FeII blend. This effect in a typically low luminosity sample, where HeII is stronger than average, likely drove the conclusion that FeII was systematically redshifted. We tested this conclusion omitting the HeII line from our fits to the bin A2 and B1 composites generated from the H08 sample. FeII shifts in lines 2 and 5 of table 1 increase from -60 to $+770 \text{ km/s}$ and from 730 to 1570 km/s , respectively. The more constraining A2 results suggest that HeII can produce the entire systematic redshift claimed by H08.

4. Conclusions

We *do not* confirm large FeII redshifts relative to narrow [OIII] and broad H β emission in Type 1 AGN but cannot rule out the existence of small red (or blue) shifts in particular subsamples. Fitting median composites built from spectra with large claimed FeII shifts ($\geq 1000 \text{ km s}^{-1}$) indicates small shifts with an upper limit $\approx 300 \text{ km s}^{-1}$ for bin A2. In the case of B1 the best fit suggests $\approx 700 \text{ km s}^{-1}$ but the shift is very poorly constrained. In both cases the shifts are not significantly different from 0. These results do not support the origin of FeII emission from a dynamical disjoint region from the one emitting the broad core of H β . Our result also challenges the usefulness of FeII shift as orientation parameter. Small systematic shifts of FeII with respect to the rest frame seem plausible but a reliable analysis is possible only on spectra of high s/n ratio.

REFERENCES

- Bachev, R., Marziani, P., Sulentic, J. W., Zamanov, R., Calvani, M., & Dultzin-Hacyan, D. 2004, *ApJ*, 617, 171
- Bevington, P. R. 1969, *Data reduction and error analysis for the physical sciences* (New York: McGraw-Hill)
- Boroson, T. 2005, *AJ*, 130, 381
- Boroson, T. A. 2011, *ApJ*, 735, L14+
- Boroson, T. A., & Green, R. F. 1992, *ApJS*, 80, 109
- Ferland, G. J., Hu, C., Wang, J., Baldwin, J. A., Porter, R. L., van Hoof, P. A. M., & Williams, R. J. R. 2009, *ApJ*, 707, L82
- Gaskell, C. M. 1982, *ApJ*, 263, 79
- . 1985, *Nature*, 315, 386
- Hu, C., Wang, J.-M., Ho, L. C., Chen, Y.-M., Bian, W.-H., & Xue, S.-J. 2008, *ApJ*, 687, 78 (H08)
- Kovačević, J., Popović, L. Č., & Dimitrijević, M. S. 2010, *ApJS*, 189, 15
- Kriss, G. 1994, *Astronomical Data Analysis Software and Systems III*, A.S.P. Conference Series, 61, 437

- Marziani, P., Sulentic, J. W., Calvani, M., Perez, E., Moles, M., & Penston, M. V. 1993, *ApJ*, 410, 56
- Marziani, P., Sulentic, J. W., Negrete, C. A., Dultzin, D., Zamfir, S., & Bachev, R. 2010, *MNRAS*, 409, 1033
- Marziani, P., Sulentic, J. W., Stirpe, G. M., Zamfir, S., & Calvani, M. 2009, *A&Ap*, 495, 83
- Marziani, P., Sulentic, J. W., Zamanov, R., Calvani, M., Dultzin-Hacyan, D., Bachev, R., & Zwitter, T. 2003a, *ApJS*, 145, 199
- Marziani, P., Zamanov, R. K., Sulentic, J. W., & Calvani, M. 2003b, *MNRAS*, 345, 1133
- Richards, G. T., et al. 2011, *AJ*, 141, 167
- Shen, Y. et al. 2011, *ApJS*, 194, 45
- Shields. G., et al 2010, *ApJ*, 687, 78
- Sulentic, J. W., Bachev, R., Marziani, P., Negrete, C. A., & Dultzin, D. 2007, *ApJ*, 666, 757
- Sulentic, J. W., Marziani, P., & Dultzin-Hacyan, D. 2000a, *ARA&A*, 38, 521
- Sulentic, J. W., Marziani, P., Zamanov, R., Bachev, R., Calvani, M., & Dultzin-Hacyan, D. 2002, *ApJL*, 566, L71
- Sulentic, J. W., Marziani, P., Zwitter, T., Dultzin-Hacyan, D., & Calvani, M. 2000b, *ApJL*, 545, L15
- Sulentic, J. W., Repetto, P., Stirpe, G. M., Marziani, P., Dultzin-Hacyan, D., & Calvani, M. 2006, *A&Ap*, 456, 929
- Véron-Cetty, M.-P., Véron, P., & Gonçalves, A. C. 2001, *AAp*, 372, 730
- Wang, T. -G., Dong, X. -B., Zhang, X., -G., Zhou, H., -Y., Wang, J. -X. & Lu, Y., -J. 2005, *ApJL*, 625, L35
- Wang, H., Wang, T., Zhou, H., Liu, B., Wang, J., Yuan, W., & Dong, X. 2011, *ArXiv e-prints*
- Zamanov, R., Marziani, P., Sulentic, J. W., Calvani, M., Dultzin-Hacyan, D., & Bachev, R. 2002, *ApJL*, 576, L9
- Zamfir, S., Sulentic, J. W., & Marziani, P., 2008, *MNRAS*, 403, 1759

Zamfir, S., Sulentic, J. W., Marziani, P., & Dultzin, D. 2010, MNRAS, 387, 856 (Z10)

Zhou, H., Wang, T., Yuan, W., Lu, H., Dong, X., Wang, J., & Lu, Y. 2006, ApJS, 166, 128

Table 1. FeII and H β shifts^a

Spectrum	$s \pm \delta s(\text{FeII})$	$s(\text{H}\beta)^{\text{b}}$ BC	$s(\text{H}\beta)$ core	$ \Delta ^{\text{c}}$	N spectra
A2 - Z10	-40 ± 90	10	0	$\lesssim 20$	130
A2 - H08, $s \geq 1000$	-60 ± 400	80	90	70	156
A2 - H08, $\eta \leq 0.1$	-160 ± 375	100	10	45	194
B1 - Z10	-340 ± 400	-150	-80	$\lesssim 20$	131
B1 - H08, $s \geq 1000$	$+730^{+400}_{-800}$	0	100	$\lesssim 20$	240
B1 - H08, $\eta \leq 0.1$	$+180 \pm 450$	-80	40	$\lesssim 20$	410
B1 - H08 $\cap$ Z10, $\eta \leq 0.1$	-150 ± 470	-170	-140	45	22

^aAll shifts in km s^{-1} with respect to average radial velocity of peaks of H β narrow component and of [OIII] λ 5007.

^b 2σ uncertainties of the H β BC measurements are $\pm 60 \text{ km s}^{-1}$ and $\pm 100 \text{ km s}^{-1}$ for pop. A and B respectively.

^cAbsolute value of peak radial velocity difference H β -[OIII]; Δ is always positive if larger than 20 km s^{-1} .

Note. — Median composites listed in the first column: A2 - Z10, B1 - Z10: A2 and B1 sources in the Z10 sample; A2 and B1 - H08 $s \geq 1000$: H08 A2 and B1 sources with reported FeII shift $s \geq 1000 \text{ km s}^{-1}$; A2 and B1 - H08 $\eta \leq 0.1$: H08 A2 and B1 sources with Eddington ration $\eta \leq 0.1$ (as estimated by H08). The last column indicates the numbers of spectra combined in the median composites. Z10-based composites have $s/n > 100$, while lowest s/n H08-based composites have $s/n > 50$. For the H08 subsamples we used spectra from the SDSS quasar catalog of Shen et al. (2011)




Fig. 1.— Fits of the $H\beta$ spectral range (left and middle panels) and normalized χ^2_v (right panels) as a function of Δv_r of the FeII template. Top row: A2 median composite of spectra with reported FeII shift $\geq 1000 \text{ km s}^{-1}$; bottom row: same for a composite of spectral type B1. Thick black line: broad component of $H\beta$; thick red line: very broad component of $H\beta$; green line: FeII template; yellow lines: narrow lines ([OIII] $\lambda 5007$ and $H\beta$ narrow component). On the right panel, black squares and gray spots indicate the normalized χ^2_v computed on the composite with no restriction on FeII width, and with FeII width restricted to roughly match the distribution of $H\beta$ width. Dot-dashed lines trace the minimum χ^2_v ratio indicating a significant difference from the best fit case. Circled symbols identify the SPECFIT best fit. The circled cross marks the best fit assuming no HeII 4686 contribution. Composites with no FeII width restriction have higher s/n making it possible to obtain more stringent limits on the maximum FeII shift.